



Non Ruminant Research Institute
P.O. Box 25-20117, Tel: + 254 708620095/7, Naivasha
E-mail: kalro.poultry@kalro.org

Hatching Date: _____

Batch No. _____

RECOMMENDED INDIGENOUS CHICKEN VACCINATION PROGRAM

AGE	VACCINE	MODE OF ADMINISTRATION	REMARKS
Day old	Mareks	Subcutaneous	Mainly for commercial hatcheries
	Newcastle+Infectious Bronchitis	Aerosol Spray	
DAY 14	Gumboro (1 st dose)	Drinking water	
Day 21	Newcastle disease (1 st dose)	Eye drop or Drinking water	
Day 24	Gumboro (2 nd dose)	Drinking water	
3 Weeks (in hot spot areas)	Fowl pox	Wing web stab	
6 Weeks (Other areas)			
8 Weeks	Newcastle disease (2 nd dose)	Eye drop or Drinking water	
	Fowl typhoid	Intramuscular injection	
18 Weeks	Newcastle disease (3 rd dose at point of lay)	Eye drop or Drinking water	Repeat every 3 months
19 Weeks	De-worming	Drinking water	Repeat every 3 months

Notes:

- NEVER vaccinate sick chicken
- Consult your veterinary/livestock staff for detailed vaccination programs in your area

