




Red spider mite on tomato

Tetranychus spp.; local name: Utitiri wekundu (Swahili)

	Prevention	Monitoring	Direct Control	Direct Control	Restrictions
 <p>Extensive webbing on a tomato plant in a greenhouse. (Photo: RSM project, ICIPE)</p>	<ul style="list-style-type: none"> ◆ Avoid water stress. If necessary, irrigate the crop regularly, and apply mulch to reduce water evaporation. ◆ Avoid continuous cropping of tomato and solanaceous plants (e.g. eggplant, capsicum and Irish potato) ◆ Practise good field sanitation e.g. removing residues of the previous crop, removing alternative hosts e.g. weeds. 	<ul style="list-style-type: none"> ◆ Inspect the crop regularly to determine the presence and level of infestation during the growing period ◆ Assess leaves particularly the lower leaves (underside) and the main vein 	<ul style="list-style-type: none"> ◆ Overhead irrigation reduces the RSM population. ◆ Use neem (Azadiractin) three to four times per season starting to spray 15 days after transplanting (2.5 to 3l per Ha) 	<ul style="list-style-type: none"> ◆ Abamectin 	<ul style="list-style-type: none"> ◆ Toxic aquatic organisms; avoid applying near water ways ◆ High risk to bees. Don't apply during flowering
	<ul style="list-style-type: none"> ◆ Clean stakes and twines with water and soap and dry in the sun for one week before transferring to a new crop. ◆ Interplanting tomato with garlic, basil or onion (repellents). 	<ul style="list-style-type: none"> ◆ Inspect whorl leaves for spots and fine webbing; Brown or silvery colour as a result of RSM feeding; and white speckling on the fruits 	<ul style="list-style-type: none"> ◆ Spray 2% horticultural oil (85ml in 4 l of water) in a six week interval 	<ul style="list-style-type: none"> ◆ Bifenthrin 	<ul style="list-style-type: none"> ◆ Moderately hazardous (WHO II) ◆ Highly toxic to bees ◆ Toxic to aquatic organisms
	<ul style="list-style-type: none"> ◆ Spray water regularly on plants to reduce dust since RSM does well in dusty conditions. ◆ Avoid planting new crops next to an already infested field (tomato or other crops) ◆ Plant tolerant varieties e.g. Rossol ◆ Intercrop with pigeon peas and spider plant, host plants for the predatory mites of spider mites. 	<ul style="list-style-type: none"> ◆ Randomly select 20 tomato plants and assess the level of mite damage of three leaflets per plant using a leaf index ranking from 1 – 5 (1 is 1-5% yellow spots on the leaf, 5 is 75-100%) ◆ Take action when the average exceeds scale 1. 	<ul style="list-style-type: none"> ◆ Garlic oil spray (85g fine chopped garlic soaked one day in 50 mls mineral oil plus 10 mls soap, stir and add 1 l water) spray twice per season. ◆ Biological control – use PHYTOTECH (<i>Phytoseiulus persimilis</i>) and AMBLYTECH (<i>Amblyseius californicus</i>) from DUDUTECH 	<ul style="list-style-type: none"> ◆ Sulphur 	<ul style="list-style-type: none"> ◆ Slightly hazardous (WHO III) ◆ Broad spectrum acaricide ◆ Sulphur is attractive to livestock hence keep livestock away
<p>Top – Red spider mite Bottom – Damaged leaf [courtesy of NRI, UK – Handbook for extension staff, Zimbabwe]</p>				<ul style="list-style-type: none"> ◆ Amitraz 	<ul style="list-style-type: none"> ◆ Moderately hazardous (WHO II) ◆ Spray only once on the underside of the leaves especially after staking and pruning.
				<ul style="list-style-type: none"> ◆ Carbamate 	<ul style="list-style-type: none"> ◆ Not classified by the WHO



Kenya

CREATED: July 2013

AUTHOR/S: Miriam Otipa (KALRO), Nancy Murage (MOAL&F), Dora Kilalo Dr. (UON), Alfayo Ombuya (KEPHIS), Willis Ochilo (CABI)

EDITED BY: Melanie Bateman (CABI) and Erica Chernoh (CABI)

LOSE LESS, FEED MORE

Plantwise is a CABI-led global initiative. www.plantwise.org